

Hemodynamic parameters to guide fluid therapy

PAUL E. MARIK, MD, FCCM, FCCP

Department of Medicine, Division of Pulmonary and Critical Care Medicine, Eastern Virginia Medical School, Norfolk, VA, USA

Correspondence to:

Dr Paul Marik, Eastern Virginia Medical School, 825 Fairfax Avenue, Suite 410, Norfolk, VA 23507, USA
E-mail: marikpe@evms.edu

Publication data

Received: 27 April 2010

Revision received: 29 May 2010

Accepted: 5 June 2010

Keywords

- Critical care
- CVP
- Echocardiography
- Fluid responsiveness
- Fluid therapy
- Heart–lung interactions
- Hemodynamic monitoring
- ICU
- Operating room
- PAOP
- Preload
- Pulse contour analysis
- Pulse pressure variation
- Stroke volume
- Stroke volume variation

The author has no financial interest in any of the products mentioned in this paper.

SUMMARY

The clinical determination of the intravascular volume can be extremely difficult in critically ill and injured patients as well as those undergoing major surgery. This is problematic as fluid loading is considered the first step in the resuscitation of hemodynamically unstable patients. Yet, multiple studies have demonstrated that only about 50% of hemodynamically unstable patients in the ICU and operating room respond to a fluid challenge. Cardiac filling pressures including the central venous pressure and pulmonary artery occlusion pressure have traditionally been used to guide fluid management. However, studies performed over the last 30 years have demonstrated that cardiac filling pressures are unable to predict fluid responsiveness. Over the last decade a number of studies have been reported that have used heart–lung interactions during mechanical ventilation to assess fluid responsiveness. Specifically, the pulse pressure variation derived from analysis of the arterial waveform, the stroke volume variation derived from pulse contour analysis and the variation of the amplitude of the pulse oximeter plethysmographic waveform have been shown to be highly predictive of fluid responsiveness. While the left ventricular end-diastolic area as determined by transesophageal echocardiography is a more accurate measure of preload than either the central venous pressure or pulmonary artery occlusion pressure, it does not predict fluid responsiveness as well as the dynamic indices. This paper reviews the evolution and accuracy of methods for assessing fluid responsiveness in patients in the ICU and operating room.

INTRODUCTION

Despite improvements in resuscitation and supportive care, progressive organ dysfunction occurs in a large proportion of patients with acute, life-threatening illnesses and those undergoing major surgery. It has been proposed that the multi-organ dysfunction syndrome of the critically ill is a consequence of tissue hypoxia due to inadequate oxygen delivery, often exacerbated by a microcirculatory injury and increased tissue metabolic demands.^{1,2} This may be further compounded by cytopathic hypoxia due to mitochondrial dysfunction.^{3,4} Emerging data suggest that early aggressive resuscitation of critically ill patients may limit and/or reverse tissue hypoxia, progression to organ failure and improve outcome.⁵ In a landmark study, Rivers *et al.* demonstrated that a protocol of early goal-directed therapy reduces organ failure and improves survival in patients with severe sepsis and septic shock.⁶ Similarly, optimization of cardiac output in patients undergoing major surgery may reduce postoperative complications and length of stay.⁷ Fluid therapy is considered the first step in the resuscitation of most patients with hypotension and shock. Uncorrected hypovolemia, leading to inappropriate infusions of vasopressor agents, may increase organ hypoperfusion and ischemia.⁸ However, overzealous fluid resuscitation has been associated with increased complications, increased length of ICU and hospital stay and increased mortality.^{9,10} The first step in the hemodynamic management of critically ill patients is to determine the adequacy of tissue/organ perfusion. While the signs of shock may be obvious those of subclinical hypoperfusion may be more subtle (see Table 1). It should however be noted, that increasing cardiac output/oxygen delivery in patients with adequate organ perfusion serves no useful purpose. Indeed, studies of yesteryear have demonstrated that targeting ‘supra-normal’ hemodynamic parameters may be harmful.^{11,12} In patients with indices of inadequate tissue perfusion fluid resuscitation is generally regarded as the first step in resuscitation. However, clinical studies have consistently demonstrated that only about 50% of hemodynamically unstable critically ill patients are volume responsive.¹³ The resuscitation of the critically ill patient therefore requires an accurate assessment of the patients’ intravascular volume status (cardiac preload) and the ability to predict the hemodynamic response following a fluid challenge (volume responsiveness).

Table 1. Clinical indices of the adequacy of tissue/organ perfusion

- Mean arterial pressure
 - Cerebral and abdominal perfusion pressures
- Urine output
- Mentation
- Capillary refill
- Skin perfusion/mottling
- Cold extremities (and cold knees)
- Blood lactate
- Arterial pH, BE and HCO₃
- Mixed venous oxygen saturation SmvO₂ (or ScvO₂)
- Mixed venous pCO₂
- Tissue pCO₂
- Skeletal muscle tissue oxygenation (StO₂)

A reduction in intravascular volume results in a fall in stroke volume, which is initially compensated for by an increase in heart rate thereby maintaining cardiac output. However, with further volume depletion cardiac output and then blood pressure falls. This is associated with a reduction in organ perfusion. At the organ level local autoregulatory mechanism comes into play in an attempt to maintain tissue perfusion. A reduction in renal perfusion normally results in dilatation of the glomerular afferent arteriole and constriction of the glomerular efferent arteriole so that glomerular capillary hydrostatic pressure and glomerular filtration rate remains constant. However, a decrease in renal perfusion pressure below the autoregulatory range (mean arterial pressure < 70 mmHg) leads to an abrupt fall in glomerular filtration rate and urine output (oliguria). In the elderly and in patients with diseases affecting the integrity of the afferent arterioles, lesser degrees of hypotension may cause a decline in renal function and oliguria. While primary renal diseases and urinary tract obstruction may lead to oliguria, intravascular volume depletion with renal hypoperfusion is the commonest cause of oliguria in clinical practice. Other features of intravascular volume depletion include a concentrated urine, postural hypotension, tachycardia and cold extremities (see Table 1).

Preload is best defined as left ventricular end-diastolic volume (LVEDV). According to the Frank–Starling principle as the preload increases left ventricular stroke volume increases until the optimal preload is achieved at which point the stroke volume remains relatively




Figure 1. The relationship between preload and stroke volume (Frank-Starling principle).

constant (see Figure 1). This optimal preload is related to the maximal overlap of the actin-myosin myofibrils. It is important to note that in an intact heart the actin-myosin links cannot be disengaged and hence there is *no descending limb* of the Frank-Starling curve. Once the left ventricle is functioning near the 'flat' part of the Frank-Starling curve fluid loading has little effect on cardiac output and only serves to increase tissue edema and to promote tissue hypoxia. In normal physiologic conditions, both ventricles operate on the ascending portion of the Frank-Starling curve.¹⁴ This mechanism provides a functional reserve to the heart in situations of acute stress. In normal individuals, an increase in preload (with volume challenge) results in a significant increase in stroke volume.¹⁵ In contrast, only about 50% of patients with circulatory failure will respond to a fluid challenge.^{13,16} Furthermore, as a result of altered left ventricular compliance and function, the position of an acutely ill patient on his/her Frank-Starling curve cannot be predicted from their preload (LVEDV) alone. In critically ill patients it is therefore important not only to determine the patients' preload (LVEDV) but their fluid responsiveness, i.e. to whether the patient will increase his/her stroke volume or cardiac output with fluid loading (i.e. have recruitable cardiac output). Simultaneously, it is important to determine the

patients' overall fluid balance and more specifically the interstitial fluid volume (third space volume). In patients with increased interstitial fluid volume it may be more appropriate to increase cardiac output by using a vasoactive agent rather than with fluid boluses alone (see Measures of volume overload).

'STATIC' MEASURES OF INTRAVASCULAR VOLUME

The central venous pressure

The central venous pressure (CVP) is frequently used to guide fluid management. Indeed, two surveys of European intensivists/anesthesiologists reported that over 90% used the CVP to guide fluid management.^{17,18} A recent Canadian survey reported that 90% of intensivists use the CVP to monitor fluid resuscitation in patients with septic shock.¹⁹ The basis for using the CVP to guide fluid management comes from the dogma that the CVP reflects intravascular volume; specifically it is widely believed that patients with a low CVP are volume depleted while patients with a high CVP are volume overloaded. Furthermore, the '5-2' rule that was popularized in the 1970's is still widely used today for guiding fluid therapy.²⁰ According to this rule, the

change in CVP following a fluid challenge is used to guide subsequent fluid management decisions.

The CVP describes the pressure of blood in the thoracic vena cava near the right atrium of the heart. The CVP is a good approximation of right atrial pressure, which is a major determinant of right ventricular filling. It has therefore been assumed that the CVP is a good indicator of right ventricular preload. Furthermore, as right ventricular stroke volume determines left ventricular filling, the CVP is assumed to be an indirect measure of left ventricular preload. However, because of the changes in venous tone, intrathoracic pressures (positive end expiratory pressure, etc.), left and right ventricular compliance and geometry that occur in critically ill patients, there is a poor relationship between the CVP and right ventricular end-diastolic volume. Furthermore, the right ventricular end-diastolic volume may not reflect the patients' position on the Frank–Starling curve and therefore his/her preload reserve.

We performed a systematic review to assess the value of the CVP in directing fluid management.²¹ We identified five studies that compared the CVP with the measured circulating blood volume while 19 studies determined the relationship between the CVP/delta-CVP and the change in cardiac performance following a fluid challenge. The pooled correlation coefficient between the CVP and the measured blood volume was 0.16 (95% CI 0.03–0.28). The pooled correlation coefficient between the baseline CVP and change in stroke index/cardiac index was 0.18 (95% CI 0.08–0.28). The pooled area under the receiver operator characteristic (ROC) curve was 0.56 (95% CI 0.51–0.61). The pooled correlation between the delta-CVP and the change in stroke index/cardiac index was 0.11 (95% CI 0.015–0.21). The results of this systematic review clearly demonstrate that there is no association between the CVP and circulating blood volume, that the CVP is a poor indicator of left and right ventricular preload and that the CVP does not predict fluid responsiveness. Based on these results we believe that the CVP should no longer be routinely measured in the ICU, operating room or emergency room.

Pulmonary artery occlusion pressure

Since the introduction of the pulmonary artery catheter almost 30 years ago the pulmonary artery occlusion pressure (PAOP) was assumed to be a reliable and valid indicator of left ventricular preload. However, it was not

long after the introduction of the pulmonary artery catheter that studies began to appear demonstrating that the PAOP was a poor reflection of preload. Recent studies have clearly demonstrated that the PAOP is a poor predictor of preload and volume responsiveness.^{16,21–23} The PAOP suffers many of the limitations of the CVP. The PAOP is a measure of left ventricular end-diastolic pressure and not LVEDV or LV preload. The use of the PAOP to measure left ventricular preload assumes a direct relationship between the left ventricular end-diastolic pressure and LVEDV. This pressure–volume curve, which describes left ventricular compliance, is normally curvilinear. Furthermore, alterations in left ventricular compliance shift the pressure–volume curve. Factors that alter left ventricular compliance include left ventricular preload, left ventricular afterload, left ventricular mass and ventricular fiber stiffness. Myocardial ischemia, sepsis, diabetes, obesity, aging, sustained tachycardia, dialysis, cardioplegia as well as other factors alter myocardial fiber stiffness. In addition, the left ventricular pressure–volume curve is affected by the degree of right ventricular filling. As the two ventricles are physically coupled by the interventricular septum and by the constraining effects of the pericardium, the end-diastolic pressure–volume curve of either ventricle is dependent upon the diastolic volume of the other. Furthermore, the PAOP is influenced by the juxtacardiac pressure, particularly if positive end expiratory pressure is used.

Left ventricular end-diastolic area

Transesophageal echocardiography (transgastric, mid-papillary short axis view) has been used to assess left ventricular dimensions in patients undergoing mechanical ventilation. The left ventricular end-diastolic area (LVEDA) has been shown to correlate well with the intrathoracic blood volume (ITBV) and global end-diastolic volume (GEDV),^{24,25} as well as with LVEDV as measured by scintigraphy.^{26–28} An end-diastolic diameter of < 25 mm and a LVEDA of < 55 cm² have been used to diagnose hypovolemia.²⁹ While a number of studies have found the LVEDA to be a good predictor of fluid responsiveness^{30–34} other studies have failed to replicate this finding.^{35–40} It should be recognized that a small LVEDA does not always reflect decreased intravascular volume. Small LV volumes can be seen with restriction to filling because of decreased ventricular

compliance (hypertrophy, ischemia), acute cor pulmonale [acute right ventricle (RV) dysfunction] and pericardial disease. Therefore, while the LVEDA may be an accurate measure of preload, preload does not necessarily translate into preload responsiveness. In addition to a decreased LVEDA, systolic obliteration of the LV cavity has been used as a sign of decreased preload.²⁹ However, LV end-systolic cavity obliteration does not necessarily imply decreased left ventricular filling.⁴¹ A major limitation of echocardiography is that it provides a snapshot of ventricular function at a single period in time. Recently, a disposable transesophageal echocardiography probe that allows continuous monitoring of LV function has been developed (ClariTEE, ImaCor, Uniondale, NY, USA). Such technology allows monitoring of LV volumes and function over time, allowing the clinician to determine the response to various therapeutic interventions.

Inferior vena caval diameter

The diameter of the inferior vena cava (IVC) as it enters the right atrium can be measured by subcostal echocardiography. A collapsed IVC is assumed to be indicative of volume depletion while a distended IVC is reflective of a high right atrial pressure. A number of authors have demonstrated that the mean end-diastolic IVC dimension correlates with mean right atrial pressure in both spontaneously breathing and mechanically ventilated patients.^{42,43} Measurement of the IVC diameter is therefore an indirect indicator of the CVP and is associated with all the limitations of CVP measurement.

ITBV index and GEDV index

Transpulmonary thermodilution using a single-indicator (cold bolus) is a minimally invasive technique that allows for the computation of the cardiac output (PiCCO monitoring system, Pulsion Medical Systems, Munich, Germany).⁴⁴ Transpulmonary thermodilution requires the use of a specific thermodilution tipped arterial catheter (usually placed in the femoral artery), which measure the change in temperature following the injection of a bolus of cold saline through a central vein (in the neck). Mathematical analysis of the transpulmonary thermodilution curve allows the calculation of the ITBV index as well as the volume of blood contained in the

four chambers of the heart, called the GEDV index (GEDVI). While the GEDVI provides a good estimate of intravascular volume and preload; it has the same limitations as the LVEDA (as measured by transesophageal echocardiography) in predicting volume responsiveness.^{32,38,45}

DYNAMIC INDICES OF INTRAVASCULAR VOLUME

Over the last decade a number of studies have been reported that have used heart–lung interactions during mechanical ventilation to assess fluid responsiveness. Specifically, the pulse pressure variation (PPV) derived from analysis of the arterial waveform, the stroke volume variation (SVV) derived from pulse contour analysis and the variation of the amplitude of the pulse oximeter plethysmographic waveform have been shown to be highly predictive of fluid responsiveness.

SVV and PPV

The principles underling the PPV (and SVV) are based on simple physiology (see Figure 2). Intermittent positive pressure ventilation induces cyclic changes in the loading conditions of the left and right ventricles. Mechanical insufflation decreases preload and increases afterload of the RV. The RV preload reduction is due to the decrease in the venous return pressure gradient that is related in the inspiratory increase in pleural pressure.⁴⁶ The increase in RV afterload is related to the inspiratory increase in transpulmonary pressure. The reduction in RV preload and increase in RV afterload both lead to a decrease in RV stroke volume, which is at a minimum at the end of the inspiratory period.⁴⁷ The inspiratory reduction in RV ejection leads to a decrease in LV filling after a phase lag of two or three heart beats because of the long blood pulmonary transit time. Thus the LV preload reduction may induce a decrease in LV stroke volume, which is at its minimum during the expiratory period. The cyclic changes in RV and LV stroke volume are greater when the ventricles operate on the steep rather than the flat portion of the Frank–Starling curve (see Figures 1 and 3). Therefore, the magnitude of the respiratory changes in LV stroke volume is an indicator of biventricular preload dependence.⁴⁶

We performed a systematic review to determine the accuracy of the PPV and SVV to predict volume respon-




Figure 2. Hemodynamic effects of mechanical ventilation. The cyclic changes in left ventricle (LV) stroke volume are mainly related to the expiratory decrease in LV preload because of the inspiratory decrease in right ventricle (RV) filling. Reproduced with permission from Critical Care/Current Science Ltd.⁴⁶




Figure 3. Arterial waveform analysis during positive pressure ventilation predicts an individual patients' position on their Starling curve and allows optimization of cardiac performance. SVV, stroke volume variation.

which the stroke volume is likely to increase. The sensitivity, specificity and diagnostic odds ratio were 0.89, 0.88 and 59.86 for the PPV and 0.82, 0.86 and 27.34 for the SVV, respectively. The predictive value was maintained in patients with poor LV function. Furthermore, with remarkable consistency these studies reported a threshold PPV/SVV of 12%–13%. In this study the area under the ROC curves were 0.55 for the CVP, 0.56 for the GEDVI and 0.64 for the LVEDA index (see Table 2). The enormous appeal of using the PPV/SVV as a marker of volume responsiveness is that it dynamically predicts an individual patients' position on their Starling curve and this is independent of ventricular function and compliance as well as pulmonary pressures and mechanics (see Figure 3). Furthermore, this technology is relatively simple both in concept and in execution and is conducive to monitoring both in the operating room and ICU. In a pilot study Lopes and colleagues demonstrated that optimization of cardiac output based on the PPV during high risk surgery significantly reduced the number of postoperative complications as well as ICU and hospital length of stay.⁷

Table 2. Predictive value of techniques used to determine fluid responsiveness¹³

Method	Technology	AUC*
Pulse pressure variation	Arterial waveform	0.94 (0.93–0.95)
Systolic pressure variation	Arterial waveform	0.86 (0.82–0.90)
Stroke volume variation	Pulse contour analysis	0.84 (0.78–0.88)
Left ventricular end-diastolic area	Echocardiography	0.64 (0.53–0.74)
Global end-diastolic volume	Transpulmonary thermodilution	0.56 (0.37–0.67)
Central venous pressure	Central venous catheter	0.55 (0.18–0.62)

*AUC = area under the curve with 95% confidence intervals.

It should be appreciated that both arrhythmias and spontaneous breathing activity will lead to misinterpretations of the respiratory variations in pulse pressure/stroke volume. Furthermore, for any specific preload condition the PPV/SVV will vary according to the tidal volume. Reuter and colleagues demonstrated a linear relationship between tidal volume and SVV.⁴⁹ De Backer and colleagues evaluated the influence of tidal volume on the ability of the PPV to predict fluid responsiveness.⁵⁰ These authors reported that the PPV was a reliable predictor of fluid responsiveness only when the tidal volume was at least 8 mL/kg. For accuracy, reproducibility and consistency we suggest that the tidal volume be increased to 8–10 mL/kg ideal body weight prior to and after a fluid challenge.

The pulse oximeter plethysmographic waveform differs from the arterial pressure waveform by measuring volume rather than pressure changes in both arterial and venous vessels. As an extension of pulse pressure analysis during mechanical ventilation, dynamic changes in both the peak and amplitude of the pulse oximeter plethysmographic waveform have been used to predict fluid responsiveness.⁵¹ The dynamic changes of the plethysmographic waveform with positive pressure ventilation have shown a significant correlation and good agreement with the PPV, and has accurately predicted fluid responsiveness in both the operating room and ICU setting.^{52–54} The ‘pleth variability index’ (PVI) is an automated measure of the dynamic change in the perfusion index that occurs during a respiratory cycle (Masimo Corporation, Irvine, CA, USA). The perfusion index is the infrared pulsatile signal indexed against the non-pulsatile signal and reflects the amplitude of the pulse oximeter waveform. The PVI correlates closely with the respiratory induced variation in the plethysmographic and arterial pressure waveforms and can predict

fluid responsiveness non-invasively in mechanically ventilated patients.^{55,56} These oximetry techniques may be valuable for monitoring fluid responsiveness in ICU and surgical patients who do not have an arterial catheter *in situ*.

Dynamic changes in aortic flow velocity/stroke volume assessed by echocardiography

The respiratory changes in aortic flow velocity and stroke volume can be assessed by Doppler echocardiography. Assuming that the aortic annulus diameter is constant over the respiratory cycle, the changes in aortic blood flow should reflect changes in LV stroke volume. Feissel and colleagues demonstrated that the respiratory changes in aortic blood velocity predicted fluid responsiveness in mechanically ventilated patients.³⁵ In this study the LVEDA index was unable to predict fluid responsiveness. Wiesenack and colleagues were however unable to replicate these findings.⁴⁵

Positive pressure ventilation induced changes in vena caval diameter

Cyclic changes in superior (SVC) and inferior vena caval (IVC) diameter as measured by echocardiography have been used to predict fluid responsiveness. Barbier and colleagues and Feissel and coworkers demonstrated that the distensibility index of the IVC, which reflects the increase in the IVC diameter on inspiration (mechanical ventilation) was able to predict fluid responsiveness.^{57,58} This technique has a number of limitations, including the fact that subcostal echocardiography may be difficult in obese patients and those that have undergone laparotomy. Furthermore, changes in IVC diameter are affected by intra-abdominal pressure (IAP) making this

technique unreliable in patients with high IAP. Vieillard-Baron and colleagues have demonstrated that the collapsibility index of the SVC is highly predictive of volume responsiveness.^{59,60} The SVC collapsibility index is considered to be more reliable than the IVC distensibility index in predicting fluid responsiveness.⁶¹ The major drawback of this technique is that the SVC can only be adequately visualized by transesophageal echocardiography and this technique is not conducive to continuous monitoring.

The dynamic indices of volume responsiveness reviewed above are dependent on the cyclic changes in intrathoracic pressure induced by positive pressure ventilation and are not applicable to spontaneously breathing patients. However, changes in aortic flow velocity and stroke volume induced by passive leg raising in non-ventilated patients have been demonstrated to be predictive of volume responsiveness.^{62,63} Similarly, Delorme and colleagues demonstrated changes in the amplitude of the pulse oximeter plethysmographic waveform during passive leg raising in spontaneously breathing individuals.⁶⁴ This technique may prove useful in assessing fluid responsiveness in spontaneously breathing patients.

MEASURES OF VOLUME OVERLOAD

Recent data strongly suggest that in critically ill and injured patients excessive fluid resuscitation with an accumulating positive fluid balance is associated with worse clinical outcomes.⁹ Data from the ARDSNet study group indicate that the cumulative fluid balance by the fourth day of study was highly predictive of hospital mortality as well as ventilator- and ICU-free days.⁶⁵ In a cohort of patients with acute lung injury secondary to septic shock Murphy and colleagues demonstrated that a fluid management strategy that included adequate initial fluid resuscitation (within the first 6 hours) followed by a conservative late fluid management strategy (even-to negative fluid balance on at least 2 days during the first 7 days of treatment) was associated with improved patients outcomes.⁶⁶

Dynamic changes in pulse pressure, stroke volume or perfusion index (PVI) together with echocardiographic assessment of LV function and size provide useful information for assessing preload and preload responsiveness; however, these techniques do not allow for the detection and quantification of fluid excess. Similarly,

the CVP and PAOP are poor measures of volume status and cannot be used reliably to detect volume overload.²¹ Some have suggested that patients receive volume resuscitation until they develop pulmonary edema (indicating that the 'tank is full'); this is clearly a dangerous approach.⁶⁷ Furthermore, radiographic and clinical signs of pulmonary edema and clinical evidence of anasarca are late signs of volume overload and poor end points for fluid resuscitation. Extravascular lung water (EVLW) as determined by transpulmonary thermodilution and IAP monitoring are two techniques that 'measure' interstitial fluid volume (tissue edema) and may aid in the assessment of volume overload.

EVLW

EVLW may be calculated from the descending limb (indicator dissipation) of the transpulmonary thermodilution curve.⁶⁸ This technique has been shown to compare favorably with the double indicator dilution technique and the ex-vivo gravimetric method.⁶⁹⁻⁷² Furthermore, this technique can detect small (10%-20%) increases in lung water.⁷³ The 'normal' value for EVLW is reported to be 5-7 mL/kg with values as high as 30 mL/kg during severe pulmonary edema. In an intriguing study, Sakka *et al.* found that the mortality was approximately 65% in ICU patients with an EVLW > 15 mL/kg whereas the mortality was 33% in patients with an EVLW < 10 mL/kg.⁷⁴ EVLW has been demonstrated to be an accurate indicator of the severity of lung injury and a reliable prognostic indicator in patients with sepsis-induced acute lung injury.^{75,76} EVLW should be indexed to ideal body weight rather than actual body weight.⁷⁷ It is likely that clinicians using EVLW measurements to guide fluid therapy may reduce the cumulative positive fluid balance, the duration of mechanical ventilation and ultimately patient outcome. Furthermore, by dynamically following a patient's preload and preload responsiveness together with his/her EVLW, will allow the clinician to determine the patient's optimal preload (see Figure 4).

IAP monitoring

IAP is the pressure concealed within the abdominal cavity.⁷⁸ The World Society of the Abdominal Compartment Syndrome (<http://www.wsacs.org>) has recently developed consensus definitions outlining standards for




Figure 4. Optimization of preload based on dynamic changes in preload responsiveness and extravascular lung water (reproduced with permission from Dr Charles Phillips). EVLW, extravascular lung water.

IAP measurement as well as diagnostic criteria for intra-abdominal hypertension (IAH). According to the consensus guidelines IAH is defined as an IAP ≥ 12 mmHg and abdominal compartment syndrome as an IAP above 20 mmHg with evidence of organ dysfunction/failure.⁷⁹ The abdominal perfusion pressure is a more accurate

predictor of visceral perfusion (MAP-IAP) with a target above 60 mmHg correlating with improved survival.⁷⁸ Major risk factors for IAH include abdominal surgery/trauma, high volume fluid resuscitation (> 3500 mL/24 hours), massive blood transfusion (> 10 units/24 hours), large burns, ileus, damage control laparotomy, liver failure with ascites, severe pancreatitis and liver transplantation. The IAP should be measured in all 'at risk patients' with repeated measures in those with IAH and following clinical deterioration. In patients with an increasing IAP volume resuscitation should be limited with attempts to achieve a negative fluid balance.

CONCLUSION

By virtue of its simplicity, accuracy and availability as a continuous monitoring tool, dynamic monitoring of the pulse pressure, stroke volume and pulse oximeter plethysmographic waveform would appear to be the ideal methods for the titration of fluid resuscitation in critically ill patients undergoing mechanical ventilation. Echocardiographic methods of assessing ventricular function and size complement the information obtained by these dynamic indices of fluid responsiveness. Measurement of EVLW and IAP may be useful in preventing volume overload. These data should be interpreted in the context of the patients' clinical condition as well as other parameters including the chest radiograph, PaO₂/FiO₂, urine output, renal function and cumulative fluid balance. The CVP and PAOP no longer have a place in modern hemodynamic monitoring.

REFERENCES

- Ince C, Sinaasappel M. Microcirculatory oxygenation and shunting in sepsis and shock. *Crit Care Med* 1999; 27: 1369-77.
- Beal AL, Cerra FB. Multiple organ failure syndrome in the 1990's: systemic inflammatory response and organ dysfunction. *JAMA* 1994; 271: 226-33.
- Fink MP. Bench-to-bedside review: cytopathic hypoxia. *Crit Care* 2002; 6: 491-9.
- Fink MP. Cytopathic hypoxia. Is oxygen use impaired in sepsis as a result of an acquired intrinsic derangement in cellular respiration? *Crit Care Clin* 2002; 18: 165-75.
- Levy MM, Macias WL, Russell JA, *et al.* Failure to improve during the first day of therapy is predictive of 28-day mortality in severe sepsis. *Chest* 2004; 124(Suppl.): 120S.
- Rivers E, Nguyen B, Havstad S, *et al.* Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 2001; 345: 1368-77.
- Lopes MR, Oliveira MA, Pereira VO, *et al.* Goal-directed fluid management based on pulse pressure variation monitoring during high-risk surgery: a pilot randomized controlled trial. *Crit Care* 2007; 11: R100.
- Murakawa K, Kobayashi A. Effects of vasopressors on renal tissue gas tensions during hemorrhagic shock in dogs. *Crit Care Med* 1988; 16: 789-92.
- Vincent JL, Sakr Y, Sprung CL, *et al.* Sepsis in European intensive care units: results of the SOAP study. *Crit Care Med* 2006; 34: 344-53.
- The ARDSNet Group. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med* 2006; 354: 2564-75.
- Hayes MA, Timmins AC, Yau E, *et al.* Elevation of systemic oxygen delivery in the treatment of critically ill patients. *N Engl J Med* 1994; 330: 1717-22.
- Gattinoni L, Brazzi L, Pelosi P, *et al.* A trial of goal-oriented hemodynamic therapy in critically ill patients. *N Engl J Med* 1995; 333: 1025-32.

- 13 Marik PE, Cavallazzi R, Vasu T, *et al.* Dynamic changes in arterial waveform derived variables and fluid responsiveness in mechanically ventilated patients. A systematic review of the literature. *Crit Care Med* 2009; 37: 2642-7.
- 14 Braunwald E, Sonnenblick EH, Ross J. Mechanisms of cardiac contraction and relaxation. In: Braunwald E (ed.). *Heart Disease*. W.B.Saunders Company: Philadelphia, PA, 1988, pp. 383-425.
- 15 Nixon JV, Murray RG, Leonard PD, *et al.* Effect of large variations in preload on left ventricular characteristics in normal subjects. *Circulation* 1982; 65: 698-703.
- 16 Michard F, Teboul JL. Predicting fluid responsiveness in ICU patients: a critical analysis of the evidence. *Chest* 2002; 121: 2000-8.
- 17 Boldt J, Lenz M, Kumle B, *et al.* Volume replacement strategies on intensive care units: results from a postal survey. *Intensive Care Med* 1998; 24: 147-51.
- 18 Kastrup M, Markewitz A, Spies C, *et al.* Current practice of hemodynamic monitoring and vasopressor and inotropic therapy in post-operative cardiac surgery patients in Germany: results from a postal survey. *Acta Anaesthesiol Scand* 2007; 51: 347-58.
- 19 McIntyre LA, Hebert PC, Fergusson D, *et al.* A survey of Canadian intensivists' resuscitation practices in early septic shock. *Crit Care* 2007; 11: R74.
- 20 Weil MH, Henning RJ. New concepts in the diagnosis and fluid treatment of circulatory shock. Thirteenth annual Becton, Dickinson and Company Oscar Schwidetsky Memorial Lecture. *Anesth Analg* 1979; 58: 124-32.
- 21 Marik PE, Baram M, Vahid B. Does the central venous pressure predict fluid responsiveness? A systematic review of the literature and the tale of seven mares. *Chest* 2008; 134: 172-8.
- 22 Osman D, Ridet C, Ray P, *et al.* Cardiac filling pressures are not appropriate to predict hemodynamic response to volume challenge. *Crit Care Med* 2007; 35: 64-8.
- 23 Michard F, Boussat S, Chemla D, *et al.* Relation between respiratory changes in arterial pulse pressure and fluid responsiveness in septic patients with acute circulatory failure. *Am J Respir Crit Care Med* 2000; 162: 134-8.
- 24 Hofer CK, Furrer L, Matter-Ensner S, *et al.* Volumetric preload measurement by thermodilution: a comparison with transoesophageal echocardiography. *Br J Anaesth* 2005; 94: 748-55.
- 25 Hinder F, Poelaert JI, Schmidt C, *et al.* Assessment of cardiovascular volume status by transoesophageal echocardiography and dye dilution during cardiac surgery. *Eur J Anaesthesiol* 1998; 15: 633-40.
- 26 Urbanowicz JH, Shabaan MJ, Cohen NH, *et al.* Comparison of transoesophageal echocardiographic and scintigraphic estimates of left ventricular end-diastolic volume index and ejection fraction in patients following coronary artery bypass grafting. *Anesthesiology* 1990; 72: 607-12.
- 27 Clements FM, Harpole DH, Quill T, *et al.* Estimation of left ventricular volume and ejection fraction by two-dimensional transoesophageal echocardiography: comparison of short axis imaging and simultaneous radionuclide angiography. *Br J Anaesth* 1990; 64: 331-6.
- 28 Ryan T, Burwash I, Lu J, *et al.* The agreement between ventricular volumes and ejection fraction by transoesophageal echocardiography or a combined radionuclear and thermodilution technique in patients after coronary artery surgery. *J Cardiothorac Vasc Anesth* 1996; 10: 323-8.
- 29 Subramaniam B, Talmor D. Echocardiography for management of hypotension in the intensive care unit. *Crit Care Med* 2007; 35: S401-7.
- 30 Preisman S, Kogan S, Berkenstadt H, *et al.* Predicting fluid responsiveness in patients undergoing cardiac surgery: functional haemodynamic parameters including the Respiratory Systolic Variation Test and static preload indicators. *Br J Anaesth* 2005; 95: 746-55.
- 31 Tavernier B, Makhotine O, Lebuffe G, *et al.* Systolic pressure variation as a guide to fluid therapy in patients with sepsis-induced hypotension. *Anesthesiology* 1998; 89: 1313-21.
- 32 Reuter DA, Felbinger TW, Schmidt C, *et al.* Stroke volume variations for assessment of cardiac responsiveness to volume loading in mechanically ventilated patients after cardiac surgery. *Intensive Care Med* 2002; 28: 392-8.
- 33 Charron C, Fessenmeyer C, Cosson C, *et al.* The influence of tidal volume on the dynamic variables of fluid responsiveness in critically ill patients. *Anesth Analg* 2006; 102: 1511-17.
- 34 Solus-Biguenet H, Fleyfel M, Tavernier B, *et al.* Non-invasive prediction of fluid responsiveness during major hepatic surgery. *Br J Anaesth* 2006; 97: 808-16.
- 35 Feissel M, Michard F, Mangin I, *et al.* Respiratory changes in aortic blood velocity as an indicator of fluid responsiveness in ventilated patients with septic shock. *Chest* 2001; 119: 867-73.
- 36 Tousignant CP, Walsh F, Mazer CD. The use of transoesophageal echocardiography for preload assessment in critically ill patients. *Anesth Analg* 2000; 90: 351-5.
- 37 Rex S, Brose S, Metzelder S, *et al.* Prediction of fluid responsiveness in patients during cardiac surgery. *Br J Anaesth* 2004; 93: 782-8.
- 38 Hofer CK, Muller SM, Furrer L, *et al.* Stroke volume and pulse pressure variation for prediction of fluid responsiveness in patients undergoing off-pump coronary artery bypass grafting. *Chest* 2005; 128: 848-54.
- 39 Belloni L, Pisano A, Natale A, *et al.* Assessment of fluid-responsiveness parameters for off-pump coronary artery bypass surgery: a comparison among LiDCO, transoesophageal echocardiography, and pulmonary artery catheter. *J Cardiothorac Vasc Anesth* 2008; 22: 243-8.
- 40 Lee JH, Kim JT, Yoon SZ, *et al.* Evaluation of corrected flow time in oesophageal Doppler as a predictor of fluid responsiveness. *Br J Anaesth* 2007; 99: 343-8.
- 41 Leung JM, Levine EH. Left ventricular end-systolic cavity obliteration as an estimate of intraoperative hypovolemia. *Anesthesiology* 1994; 81: 1102-9.
- 42 Mintz GS, Kotler MN, Parry WR, *et al.* Real-time inferior vena caval ultrasonography: normal and abnormal findings and its use in assessing right-heart function. *Circulation* 1981; 64: 1018-25.
- 43 Jue J, Chung W, Schiller NB. Does inferior vena cava size predict right atrial pressures in patients receiving mechanical ventilation? *J Am Soc Echocardiogr* 1992; 5: 613-19.
- 44 Genahr A, McLuckie A. Transpulmonary thermodilution in the critically ill. *Br J Intensive Care* 2004; Spring: 6-10.
- 45 Wiesenack C, Fiegl C, Keyser A, *et al.* Assessment of fluid responsiveness in mechanically ventilated cardiac surgical patients. *Eur J Anaesthesiol* 2005; 22: 658-65.

- 46 Michard F, Teboul JL. Using heart-lung interactions to assess fluid responsiveness during mechanical ventilation. *Crit Care* 2000; 4: 282-9.
- 47 Theres H, Binkau J, Laule M, et al. Phase-related changes in right ventricular cardiac output under volume-controlled mechanical ventilation with positive end-expiratory pressure. *Crit Care Med* 1999; 27: 953-8.
- 48 Marik PE, Cavallazzi R, Vasu T, et al. Stroke volume variation and fluid responsiveness. A systematic review of the literature. *Crit Care Med* 2009; 37: 2642-7.
- 49 Reuter DA, Bayerlein J, Goepfert MS, et al. Influence of tidal volume on left ventricular stroke volume variation measured by pulse contour analysis in mechanically ventilated patients. *Intensive Care Med* 2003; 29: 476-80.
- 50 De Backer D, Heenen S, Piagnerelli M, et al. Pulse pressure variations to predict fluid responsiveness: influence of tidal volume. *Intensive Care Med* 2005; 31: 517-23.
- 51 Desebbe O, Cannesson M. Using ventilation-induced plethysmographic variations to optimize patient fluid status. *Curr Opin Anaesthesiol* 2008; 21: 772-8.
- 52 Natalini G, Rosano A, Taranto M, et al. Arterial versus plethysmographic dynamic indices to test responsiveness for testing fluid administration in hypotensive patients: a clinical trial. *Anesth Analg* 2006; 103: 1478-84.
- 53 Cannesson M, Besnard C, Durand PG, et al. Relation between respiratory variations in pulse oximetry plethysmographic waveform amplitude and arterial pulse pressure in ventilated patients. *Crit Care* 2005; 9: R562-8.
- 54 Feissel M, Teboul JL, Merlani P, et al. Plethysmographic dynamic indices predict fluid responsiveness in septic ventilated patients. *Intensive Care Med* 2007; 33: 993-9.
- 55 Cannesson M, Desebbe O, Rosamel P, et al. Pleth variability index to monitor the respiratory variations in the pulse oximeter plethysmographic waveform amplitude and predict fluid responsiveness in the operating theatre. *Br J Anaesth* 2008; 101: 200-6.
- 56 Cannesson M, Delannoy B, Morand A, et al. Does the Pleth variability index indicate the respiratory-induced variation in the plethysmogram and arterial pressure waveforms? *Anesth Analg* 2008; 106: 1189-94.
- 57 Barbier C, Loubieres Y, Schmit C, et al. Respiratory changes in inferior vena cava diameter are helpful in predicting fluid responsiveness in ventilated septic patients. *Intensive Care Med* 2004; 30: 1740-6.
- 58 Feissel M, Michard F, Faller JP, et al. The respiratory variation in inferior vena cava diameter as a guide to fluid therapy. *Intensive Care Med* 2004; 30: 1834-7.
- 59 Vieillard-Baron A, Augarde R, Prin S, et al. Influence of superior vena caval zone condition on cyclic changes in right ventricular outflow during respiratory support. *Anesthesiology* 2001; 95: 1083-8.
- 60 Vieillard-Baron A, Chergui K, Rabiller A, et al. Superior vena caval collapsibility as a gauge of volume status in ventilated septic patients. *Intensive Care Med* 2004; 30: 1734-9.
- 61 Charron C, Caille V, Jardin F, et al. Echocardiographic measurement of fluid responsiveness. *Curr Opin Crit Care* 2006; 12: 249-54.
- 62 Maizel J, Airapetian N, Lorne E, et al. Diagnosis of central hypovolemia by using passive leg raising. *Intensive Care Med* 2007; 33: 1133-8.
- 63 Lamia B, Ochagavia A, Monnet X, et al. Echocardiographic prediction of volume responsiveness in critically ill patients with spontaneously breathing activity. *Intensive Care Med* 2007; 33: 1125-32.
- 64 Delorme S, Renault R, Le MY, et al. Variations in pulse oximetry plethysmographic waveform amplitude induced by passive leg raising in spontaneously breathing volunteers. *Am J Emerg Med* 2007; 25: 637-42.
- 65 Rosenberg AL, Dechert RE, Park PK, et al. Review of a large clinical series: association of cumulative fluid balance on outcome in acute lung injury: a retrospective review of the ARDSnet tidal volume study cohort. *J Intensive Care Med* 2009; 24: 35-46.
- 66 Murphy CV, Schramm GE, Doherty JA, et al. The importance of fluid management in acute lung injury secondary to septic shock. *Chest* 2009; 136: 102-9.
- 67 Hollenberg SM, Ahrens TS, Annane D, et al. Practice parameters for hemodynamic support of sepsis in adult patients: 2004 update. *Crit Care Med* 2004; 32: 1928-48.
- 68 Isakow W, Schuster DP. Extravascular lung water measurements and hemodynamic monitoring in the critically ill: bedside alternatives to the pulmonary artery catheter. *Am J Physiol Lung Cell Mol Physiol* 2006; 291: 1118-31.
- 69 Michard F, Schachtrupp A, Toens C. Factors influencing the estimation of extravascular lung water by transpulmonary thermodilution in critically ill patients. *Crit Care Med* 2005; 33: 1243-7.
- 70 Reuter DA, Felbinger TW, Moerstedt K, et al. Intrathoracic blood volume index measured by thermodilution for preload monitoring after cardiac surgery. *J Cardiothorac Vasc Anesth* 2002; 16: 191-5.
- 71 Sakka SG, Ruhl CC, Pfeiffer UJ, et al. Assessment of cardiac preload and extravascular lung water by single transpulmonary thermodilution. *Intensive Care Med* 2000; 26: 180-7.
- 72 Katzenelson R, Perel A, Berkenstadt H, et al. Accuracy of transpulmonary thermodilution versus gravimetric measurement of extravascular lung water. *Crit Care Med* 2004; 32: 1550-4.
- 73 Fernandez-Mondejar E, Rivera-Fernandez R, Garcia-Delgado M, et al. Small increases in extravascular lung water are accurately detected by transpulmonary thermodilution. *J Trauma* 2005; 59: 1420-3.
- 74 Sakka SG, Klein M, Reinhart K, et al. Prognostic value of extravascular lung water in critically ill patients. *Chest* 2002; 122: 2080-6.
- 75 Chung FT, Lin SM, Lin SY, et al. Impact of extravascular lung water index on outcomes of severe sepsis patients in a medical intensive care unit. *Respir Med* 2008; 102: 956-61.
- 76 Kuzkov VV, Kirov MY, Sovershaev MA, et al. Extravascular lung water determined with single transpulmonary thermodilution correlates with the severity of sepsis-induced acute lung injury. *Crit Care Med* 2006; 34: 1647-53.
- 77 Berkowitz DM, Danai PA, Eaton S, et al. Accurate characterization of extravascular lung water in acute respiratory distress syndrome. *Crit Care Med* 2008; 36: 1803-9.
- 78 Cheatham ML. Intraabdominal pressure monitoring during fluid resuscitation. *Curr Opin Crit Care* 2008; 14: 327-33.
- 79 Malbrain ML, Cheatham ML, Kirkpatrick A, et al. Results from the International Conference of Experts on Intra-abdominal Hypertension and Abdominal Compartment Syndrome. I. Definitions. *Intensive Care Med* 2006; 32: 1722-32.